



Generalized anxiety disorder, depressive symptoms and sleep quality during COVID-19 outbreak in China: a web-based cross-sectional survey

Yeen Huang^{a,b}, Ning Zhao^{a,b,*}

^a The 6th Affiliated Hospital of Shenzhen University Health Science Center, Nanshan Hospital Affiliated to Shenzhen University, Shenzhen, 518052, People's Republic of China

^b Department of Rehabilitation Medicine, Huazhong University of Science and Technology Union Shenzhen Hospital, Shenzhen, 518052, People's Republic of China

ARTICLE INFO

Keywords:

COVID-19
Mental health
Anxiety
Depressive symptoms
Sleep

ABSTRACT

China has been severely affected by Coronavirus Disease 2019 (COVID-19) since December, 2019. We aimed to assess the mental health burden of Chinese public during the outbreak, and to explore the potential influence factors. Using a web-based cross-sectional survey, we collected data from 7,236 self-selected volunteers assessed with demographic information, COVID-19 related knowledge, generalized anxiety disorder (GAD), depressive symptoms, and sleep quality. The overall prevalence of GAD, depressive symptoms, and sleep quality of the public were 35.1%, 20.1%, and 18.2%, respectively. Younger people reported a significantly higher prevalence of GAD and depressive symptoms than older people. Compared with other occupational group, healthcare workers were more likely to have poor sleep quality. Multivariate logistic regression showed that age (< 35 years) and time spent focusing on the COVID-19 (≥ 3 hours per day) were associated with GAD, and healthcare workers were at high risk for poor sleep quality. Our study identified a major mental health burden of the public during the COVID-19 outbreak. Younger people, people spending too much time thinking about the outbreak, and healthcare workers were at high risk of mental illness. Continuous surveillance of the psychological consequences for outbreaks should become routine as part of preparedness efforts worldwide.

1. Introduction

Coronavirus Disease 2019 (COVID-19, also known as 2019-nCoV), a cluster of acute respiratory illness with unknown causes, has occurred in Wuhan, Hubei Province, China since December 2019 (Wuhan Municipal Health Commission, 2020; Paules et al., 2020; Wang et al., 2020). As of March 25, 2020, a total of 81,846 COVID-19 cases in China have been confirmed and 3,287 Chinese died from the disease. Internationally, sporadic cases exported from Wuhan were reported in 193 countries (such as 69,176 cases in Italy, 42,058 cases in Spain, and 53,588 cases in United States), 5 continents, and 1 international conveyance (712 cases in “Diamond Princess”) (World Health Organization, 2020a). On January 23, Wuhan city closed all access routes to stop the spread of disease. Seven days later, the World Health Organization (WHO) declared the COVID-19 as a Public Health Emergency of International Concern (PHEIC) (World Health Organization, 2020b). On March 11, the WHO declared the COVID-19 outbreak could be characterized as a “pandemic” as the virus spreads increasingly worldwide.

In addition to physical damage, COVID-19 also has a serious impact on the mental health of the public. In January 20, China confirmed human-to-human transmission of COVID-19 and some medical staff in Wuhan had been infected (XINHUANET, 2020). Since then, the public has shown anxiety-related behaviors, causing a significant shortage of medical masks and alcohol across the country. On the night of January 31, due to a news that “Shuanghuanglian oral liquid” could suppress COVID-19 (People's daily of China, 2020), the public rushed to pharmacies overnight to buy this drug. In addition, many front-line medical staff work more than 16 hours a day on average, causing them to not getting enough sleep. Unfortunately, a 37-year-old Japanese government worker who was in charge of isolated returnees from Wuhan was found to have died from apparent suicide (The Japan Times, 2020).

Evidence indicated that COVID-19 is a distinct clade from the betacoronaviruses related to human severe acute respiratory syndrome (SARS) and Middle East respiratory syndrome (MERS) (Zhu et al., 2020). Several studies showed that mental health problems could occur in both healthcare workers and SARS survivors during the SARS epidemic (Lee et al., 2007; Lu et al., 2006; McAlonan et al., 2007). Post-

* Corresponding author: The 6th Affiliated Hospital of Shenzhen University Health Science Center, Nanshan Hospital Affiliated to Shenzhen University, Shenzhen, 518052, People's Republic of China.

E-mail address: zhaoning2018@email.szu.edu.cn (N. Zhao).

<https://doi.org/10.1016/j.psychres.2020.112954>

Received 20 March 2020; Received in revised form 26 March 2020; Accepted 26 March 2020

Available online 12 April 2020

0165-1781/ © 2020 Elsevier B.V. All rights reserved.

traumatic stress disorder (PTSD) and depressive disorders were the most prevalent long-term psychological condition (Mak et al., 2009). Similar results have been reported in the previous study of MERS (Lee et al., 2018). Based on the above research evidence, we have reason to speculate that the psychological condition of the public may also be affected during COVID-19 outbreak.

Therefore, using a web-based cross-sectional study, we aimed to assess the mental health burden of Chinese public during COVID-19 outbreak, and to explore the potential influence factors. We hope that our study findings will provide data support for the targeted interventions on psychological health in Chinese public during the outbreak.

2. Methods

2.1. Study design and participants

To prevent the spread of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) through droplets or contact, we used a web-based cross-sectional survey based on the National Internet Survey on Emotional and Mental Health (NISEMH), an ongoing, online health-related behavior survey of Chinese population, to collected data. This web-based survey of the COVID-19 was sent on the Internet through the WeChat public platform and the mainstream media. All Chinese people using WeChat or other social tools may see this survey, and answered the questionnaire by scanning the Quick Response code (QR code) of the questionnaire address or clicking the relevant link. To encourage the recruitment of potential participants, all participants in the survey can receive a report on their mental health after completing the evaluation. This web-based questionnaire was completely voluntary and non-commercial.

2.2. Data collection

Participants answered the questionnaires anonymously on the Internet from February 3 to, 2020 to February 17, 2020. All subjects reported their demographic data, COVID-19 related information, and completed three standardized questionnaires which assessed their generalized anxiety disorder (GAD), depressive symptoms, and sleep quality. In order to ensure the quality of survey, we set the response range of some items (e.g., the age range was limited to 6-80 years old, some items needed to be answered in reverse) and encouraged participants to answer carefully through questionnaire explanations. In addition, questionnaires that were completed < 1 minute or > 60 minutes would be excluded from analysis. Finally, a total of 7,236 participants who completed the questionnaires were included in the analysis.

2.3. Ethical statement

This study was conducted in accordance with the Declaration of Helsinki, and was approved by the Ethics Committee of Huazhong University of Science and Technology Union Shenzhen Hospital. Electronic informed consent was obtained from each participant prior to starting the investigation. Participant could withdraw from the survey at any moment without providing any justification.

2.4. Measures

2.4.1. Demographic information

Demographic variables included gender (male or female), age, and occupation. Occupation included the following four types: (1) Healthcare workers, which included doctors, nurses, and health-related administrators; (2) Enterprise or institution workers, which consisted of enterprise employees, national/provincial/municipal institution workers, and other relevant staff; (3) Teachers or students, which included teachers or students from universities, middle schools, or elementary schools; and (4) Others, which consisted of freelancers, retirees, social workers, and other relevant staff.

2.4.2. COVID-19 related knowledge

This section was evaluated by two items: (1) Time spent focusing on the COVID-19, which measured the average time spent focusing on the COVID-19 outbreak information every day; (2) Knowledge of the COVID-19, which was assessed based on the following six judgment questions about COVID-19 related knowledge: a. Inhalation of droplets from sneezing, coughing, or talking to an infected person could cause infection; b. Contact with something contaminated by an infected person could lead to infection; c. The incubation period of the virus does not exceed 14 days; d. Contact with an asymptomatic person might also lead to infection; e. There are already targeted drugs that could cure the disease; f. Taking "Shuanghuanglian Oral Liquid" could prevent infection of this disease. Of the above six questions, one point was given for each correct answer, and no points were given for each incorrect or uncertain answer. Participants with scores ≥ 5 points, equal to 4 points, and ≤ 3 points were considered to quite understand, generally understand, and do not understand.

2.4.3. Generalized anxiety disorder

We used Chinese version of GAD-7 (Generalized Anxiety Disorder-7) scale to assess subject's anxiety symptoms. The GAD-7 has been previously used in Chinese populations, and found to have good reliability (Cronbach's $\alpha = 0.90$) (Tong et al., 2016; Wang et al., 2018). Seven items assessed the frequency of anxiety symptoms over the past two weeks on a 4-point Likert-scale ranging from 0 (never) to 3 (nearly every day). The total score of GAD-7 ranged from 0 to 21, with increasing scores indicating more severe functional impairments as a result of anxiety (Spitzer et al., 2006). For the purpose of this study, we defined a GAD-total score of 9 points or greater as the presence of anxiety symptoms (Wang et al., 2018).

2.4.4. Depressive symptoms

The Center for Epidemiology Scale for Depression (CES-D) in Chinese version was used to identify whether participants had

Table 1
Demographic characteristics of study sample ($N = 7,236$).

Variable	n (%)
Total	7236 (100.0)
Gender	
Male	3284 (45.4)
Female	3952 (54.6)
Age (Mean \pm SD)	35.3 \pm 5.6
< 35 years	3155 (43.6)
≥ 35 years	4081 (56.4)
Occupation	
Healthcare workers ^a	2250 (31.1)
Enterprise or institution workers ^b	1809 (25.0)
Teachers or students ^c	1404 (19.4)
Others ^d	1773 (24.5)
Time spent focusing on the COVID-19^e	
< 1 hour	1454 (20.1)
1-2 hours	2627 (36.3)
≥ 3 hours	3155 (43.6)
Knowledge of the COVID-19	
Do not understand (score ≤ 3 points)	398 (5.5)
General understand (score 4 points)	1136 (15.7)
Quite understand (score ≥ 5 points)	5702 (78.8)

Abbreviations: n, number; SD, Standard deviation; COVID-19, 2019 Corona Virus Disease.

^a Included doctors, nurses and health administrators.

^b Included enterprise employees, national/provincial/municipal institution workers and other relevant staff.

^c Included teachers or students from universities, middle schools, or elementary schools.

^d Included freelancers, retiree, social worker and other relevant staff.

^e Average time spent focusing on the COVID-19 outbreak information every day.

Table 2

Prevalence of GAD, depressive symptoms, and sleep quality during COVID-19 outbreak in Chinese population stratified by gender (N=7,236).

Variables	Total (N = 7236) n (%)	Male (N = 3284) n (%)	Female (N = 3952) n (%)	χ^2	P-value
GAD ^a				2.89	0.089
No	4696 (64.9)	2092 (63.7)	2394 (65.9)		
Yes	2540 (35.1)	1192 (36.3)	1348 (34.1)		
Depressive symptoms ^b				3.67	0.055
No	5782 (79.9)	2625 (80.0)	3155 (79.8)		
Yes	1454 (20.1)	657 (20.0)	797 (20.2)		
Sleep quality ^c				2.59	0.108
Good	5919 (81.8)	2660 (81.0)	3259 (82.5)		
Poor	1317 (18.2)	624 (19.0)	693 (17.5)		

Abbreviations: n, number, GAD, generalized anxiety disorder.

^a GAD was defined as individuals who scored ≥ 9 points.^b Depressive symptoms included individuals who scored > 28 points.^c Poor sleep quality was defined as individuals who scored > 7 points.

depressive symptoms (Zhang et al., 2010), and the Chinese version of this scale has been validated and extensively utilized in Chinese population (Zhang et al., 2010; Zhang and Li, 2011). Twenty items assessed the frequency of depressive symptoms over the past two weeks on a 4-point Likert-scale ranging from 0 (rarely or none of the time) to 3 (most or all of the time). The score range of the CES-D was 0-60 points, and higher scores indicated more severe depressive symptomatology (Radloff, 1977). In our study, CES-D scores greater than 28 points indicated depressive symptoms.

2.4.5. Sleep quality

The Chinese version of the PSQI (Pittsburgh Sleep Quality Index) scale was used to assess the subject's sleep quality over the past two week (Liu et al., 1996). The PSQI scale contains seven components (subjective sleep quality, sleep duration, sleep latency, habitual sleep efficiency, use of sleep medications, sleep disturbance, and daytime dysfunction), and the score for each component ranges from 0 to 3 points. The global PSQI score ranges from 0 to 21, with higher scores indicating more severe sleep disorder (Buysse et al., 1989). The Chinese version of PSQI has been demonstrated to be reliable and valid in Chinese population (Liu et al., 1996), and a global PSQI score greater than 7 points indicates poor sleep quality.

2.4.6. Statistical analysis

First, descriptive analyses were conducted to describe the demographic characteristics and COVID-19 related knowledge in Chinese population. Second, the prevalence of GAD, depressive symptoms, and

sleep quality stratified by gender, age, and occupation were reported, and Chi-square test (χ^2) was used to compare the differences between groups. Third, univariate and multivariate logistic regression models were performed to explore potential influence factors for GAD, depressive symptoms, and sleep quality during COVID-19 outbreak. Odds ratio (OR), adjusted odds ratio (AOR), and 95% confidence interval (95% CI) were obtained from logistic regression models. All data were analyzed using Statistical Package for Social Sciences (SPSS) version 24.0. P-values of less than 0.05 were considered statistically significant (2-sided tests).

3. Results

3.1. Demographic characteristics

The characteristics of participants were shown in Table 1. Of the 7,236 samples analyzed, 3,284 (45.4%) were males and 3,952 (54.6%) were females, and the mean (standard deviation) age of the participants was 35.3 ± 5.6 years. Among these samples, 2,250 (31.1%) of participants were healthcare workers, 3,155 (43.6%) of participants spent 3 hours or more a day focusing on the COVID-19 outbreak, and 5,702 (78.8%) of participants quite understand knowledge of the COVID-19.

3.2. Prevalence of GAD, depressive symptoms, and sleep quality during COVID-19 outbreak stratified by gender, age, and occupations

The prevalence of GAD, depressive symptoms, and sleep quality stratified by gender, age, and occupations were shown in Table 2, Table 3, and Table 4, respectively. The overall prevalence of GAD, depressive symptoms, and sleep quality were 35.1%, 20.1%, and 18.2%, respectively. There was no statistically significant difference in the prevalence of GAD, depressive symptoms, and sleep quality by gender ($P > 0.05$, as shown in Table 2). The prevalence of GAD and depressive symptoms was significantly higher in participants younger than 35 years than in participants aged 35 years or older ($P < 0.001$, as shown in Table 3). Compared with other occupational groups, healthcare workers (23.6%) reported the highest rate of poor sleep quality ($P < 0.001$, as shown in Table 4).

3.3. Association of influence factors with GAD, depressive symptoms, and sleep quality during COVID-19 outbreak

The associations of potential influence factors with GAD, depressive symptoms, and sleep quality during COVID-19 outbreak were presented in Table 5. In the univariate logistic regression models, age (OR = 1.77, 95% CI: 1.38-1.95) and time spent focusing on the COVID-19 (OR = 1.91, 95% CI: 1.77-2.15) were significantly associated with GAD

Table 3

Prevalence of GAD, depressive symptoms, and sleep quality during COVID-19 outbreak in Chinese population stratified by age (N=7,236).

Variables	Total (N = 7236) n (%)	Age < 35 year (N = 3155) n (%)	Age \geq 35 year (N = 4081) n (%)	χ^2	P-value
GAD ^a				20.67	<0.001
No	4696 (64.9)	1956 (62.0)	2740 (67.1)		
Yes	2540 (35.1)	1199 (38.0)	1341 (32.9)		
Depressive symptoms ^b				13.91	<0.001
No	5782 (79.9)	2458 (77.9)	3324 (81.5)		
Yes	1454 (20.1)	697 (22.1)	757 (18.5)		
Sleep quality ^c				0.58	0.446
Good	5919 (81.8)	2575 (81.6)	3344 (81.9)		
Poor	1317 (18.2)	580 (18.4)	737 (18.1)		

Abbreviations: n, number, GAD, generalized anxiety disorder.

^a GAD was defined as individuals who scored ≥ 9 points.^b Depressive symptoms included individuals who scored > 28 points.^c Poor sleep quality was defined as individuals who scored > 7 points.

Table 4Prevalence of GAD, depressive symptoms, and sleep quality during COVID-19 outbreak in Chinese population stratified by Occupations ($N=7,236$).

Variables	Total ($N=7236$) n (%)	Healthcare workers ($N=2250$) n (%)	Enterprise or institution workers ($N=1809$) n (%)	Teachers or students ($N=1404$) n (%)	Others ($N=1773$) n (%)	χ^2	P-value
GAD ^a						2.36	0.501
No	4696 (64.9)	1448 (64.4)	1179 (65.2)	911 (64.9)	1158 (65.3)		
Yes	2540 (35.1)	802 (35.6)	630 (34.8)	493 (35.1)	615 (34.7)		
Depressive symptoms ^b						2.71	0.439
No	5782 (79.9)	1804 (80.2)	1445 (79.9)	1109 (79.0)	1424 (80.3)		
Yes	1454 (20.1)	446 (19.8)	364 (20.1)	295 (21.0)	349 (19.7)		
Sleep quality ^c						98.82	<0.001
Good	5919 (81.8)	1719 (76.4)	1579 (87.3)	1203 (85.7)	1418 (80.5)		
Poor	1317 (18.2)	531 (23.6)	230 (12.7)	201 (14.3)	355 (20.0)		

Abbreviations: n, number, GAD, generalized anxiety disorder.

^a GAD was defined as individuals who scored ≥ 9 points.^b Depressive symptoms included individuals who scored > 28 points.^c Poor sleep quality was defined as individuals who scored > 7 points.

in Chinese public. Similarly, age was associated with depressive symptoms ($OR=1.80$, 95% CI : 1.35-2.01), but not with sleep quality ($OR=0.69$, 95% CI : 0.35-1.05). Occupations were related to sleep quality during COVID-19 outbreak in Chinese public ($OR=1.48$, 95% CI : 1.15-1.95).

In the multivariate logistic regression models, the above associations weakened but there were still statistical difference. Participants under 35 years were more likely to have GAD than those 35 years and older ($AOR=1.65$, 95% CI : 1.49-2.02). Besides, participants who were concerned about the COVID-19 outbreak for 3 hours or more were more likely to develop GAD than those less than 1 or 2 hours ($AOR=1.83$, 95% CI : 1.53-2.19). Similarly, participants under 35 were associated with higher risk for depressive symptoms than those 35 years and older ($AOR=1.77$, 95% CI : 1.58-2.07). Compared with other occupation participants, healthcare workers were more likely to report poor sleep quality ($AOR=1.32$, 95% CI : 1.18-1.88).

4. Discussion

Our web-based study shows a high prevalence of GAD and poor sleep quality in the Chinese public during COVID-19 outbreak. Anxiety symptoms were more likely to occur in people younger than 35 years and those who spent too much time focusing on the outbreak. Compared with other professions, healthcare workers were associated with higher risk for poor sleep quality. Our findings provided data support for accurately understanding the source of public's panic during COVID-19 outbreak.

The data in this study suggested that public's levels of anxiety-related symptoms increased when a major infectious disease occurred. Similar to the psychological burden caused by SARS (Su et al., 2007), we found that 1/3 participants showed anxiety disorders, and this mood was not different between male and female during COVID-19 outbreak, which was different from previous research that women were more likely to have anxiety than men (Guo et al., 2016; Gao et al., 2020). In addition, nearly 1/5 participants had depressive symptoms and sleep

Table 5Results of univariate and multivariate logistic regression analyses ($N=7,236$).

Variables	GAD OR (95% CI)	AOR (95% CI)	Depressive symptoms OR (95% CI)	AOR (95% CI)	Sleep quality OR (95% CI)	AOR (95% CI)
Gender						
Male	1.00	1.00	1.00	1.00	1.00	1.00
Female	1.32 (0.90-1.69)	1.22 (0.86-1.64)	1.30 (0.82-2.07)	1.24 (0.77-1.99)	0.89 (0.57-1.39)	0.82 (0.52-1.29)
Age						
< 35 years	1.77 (1.38-1.95)*	1.65 (1.49-2.02)*	1.80 (1.35-2.01)*	1.77 (1.58-2.07)*	0.69 (0.35-1.05)	0.68 (0.42-1.11)
≥ 35 years	1.00	1.00	1.00	1.00	1.00	1.00
Occupations						
Healthcare workers ^a	1.30 (0.83-2.04)	1.30 (0.82-2.08)	1.15 (0.67-1.99)	1.02 (0.58-1.81)	1.48 (1.15-1.95)*	1.32 (1.18-1.88)*
Enterprise or institution workers ^b	0.85 (0.52-1.38)	0.91 (0.55-1.49)	0.80 (0.44-1.47)	0.80 (0.44-1.49)	0.60 (0.33-1.11)	0.59 (0.32-1.10)
Teachers or students ^c	1.51 (0.91-2.53)	1.41 (0.80-2.50)	1.24 (0.67-2.31)	0.94 (0.47-1.88)	0.69 (0.35-1.35)	0.87 (0.42-1.82)
Others ^d	1.00	1.00	1.00	1.00	1.00	1.00
Time spent focusing on the COVID-19 ^e						
< 1 hour	1.00	1.00	1.00	1.00	1.00	1.00
1-2 hours	0.96 (0.59-1.57)	1.01 (0.61-1.64)	0.71 (0.40-1.27)	0.74 (0.41-1.32)	0.90 (0.50-1.62)	0.81 (0.44-1.49)
≥ 3 hours	1.91 (1.77-2.15)*	1.83 (1.53-2.19)*	0.98 (0.57-1.68)	1.11 (0.63-1.93)	1.18 (0.68-2.07)	1.02 (0.57-1.82)
Knowledge of the COVID-19						
Do not understand	1.00	1.00	1.00	1.00	1.00	1.00
General understand	0.73 (0.32-1.71)	0.68 (0.29-1.60)	0.97 (0.32-2.97)	0.90 (0.29-2.76)	1.06 (0.35-3.21)	0.92 (0.30-2.82)
Quite understand	0.93 (0.45-1.93)	0.80 (0.38-1.69)	1.30 (0.49-3.47)	1.12 (0.42-3.02)	1.29 (0.48-3.42)	1.15 (0.42-3.14)

Abbreviations: GAD, generalized anxiety disorder; OR, odds ratio; AOR, adjusted odds ratio; 95% CI , 95% confidence interval; COVID-19, 2019 Corona Virus Disease.^a Included doctors, nurses and health administrators.^b Included enterprise employees, national/provincial/municipal institution workers and other relevant staff.^c Included teachers or students from universities, middle schools, or elementary schools.^d Included freelancers, retiree, social worker and other relevant staff.^e Average time spent focusing on the COVID-19 outbreak information every day.* $P < 0.001$.

problems, indicating that the uncertainty of the epidemic progression would cause greater psychological pressure on the public. The possible reason for these mental problems may be related to the “hypochondriac concerns” (worry about being infected) (Furer et al., 1997) and feared that the epidemic was hard to control.

After multivariate logistic regression analyses, we found that age and time spent focusing on COVID-19 may be potential risk factors for the psychological problems of the public. Younger participants (< 35 years) were more likely to develop anxiety and depressive symptoms during COVID-19 outbreak than older participants (≥ 35 years). Our results were similar to those of a previous study in Taiwan during SARS outbreak (Su et al., 2007). In addition, we assessed the average time participants spent focusing on the COVID-19 outbreak each day, and found that people who spent too much time thinking about the outbreak (≥ 3 hours) were more likely to develop anxiety symptoms. The manifestation of this panic mood may be related to the body's normal protective response to the stress caused by the epidemic (Maunder et al., 2003).

Since January 20, 2020, Zhong Nanshan (the renowned Chinese respiratory expert who discovered the SARS virus) confirmed that there must be human-to-human transmission of COVID-19 (XINHUA-NET, 2020), more than 40,000 medical staff have given up the Spring Festival holiday and voluntarily applied to fight against the outbreak in Hubei Province (National Health Commission of the People's Republic of China, 2020). Meanwhile, most healthcare workers in China have returned to work to cope with the further development of the disease. Our findings showed that nearly one in four healthcare workers had sleep problems, which was significantly higher than other occupational groups. One possible reason is that the working time and labour intensity of healthcare workers will increase in the face of severe epidemic (such as SARS and MERS), causing them not to have enough time to rest, and to be prone to chronic stress and psychological distress (Lu et al., 2006; Lee et al., 2018; McAlonan et al., 2007). In severe cases, a post-traumatic stress disorder (PTSD) symptoms may even occur, which is highly correlated with poor sleep (Kobayashi et al., 2007).

Fortunately, the Chinese government has taken many strong national measures in time to avoid further spread of the COVID-19 outbreak, including requiring uninfected people to isolate themselves at home, prohibiting all gathering activities, and forcing everyone to wear medical masks to enter public places. However, there is still lack of relevant research on the targeted intervention of the public's psychological problems during the COVID-19 outbreak. We filled this research gap by analyzing the prevalence of mental health burden in Chinese public stratified by demographic characteristics and exploring related influential factors. Several appropriate interventions are recommended as follows: First, psychological interventions should be directed to vulnerable populations which include the suspected and diagnosed patients, young people, and healthcare workers, especially physicians and nurses working directly with patients or quarantined people. Second, try to control and limited the time of receiving COVID-19 related information within two hours a day, focus only on the necessary information (such as facts and data) and avoid receiving too many harmful rumors (Grein et al., 2000). Third, maintain a normal rhythm of work and rest as much as possible, exercise regularly to promote sleep quality, and do not pay too much attention to outbreak information before going to sleep.

This study has several limitations. First, since the data and relevant analyses presented here were derived from a cross-sectional design, it is difficult to make causal inferences. Second, the study was limited to COVID-19 outbreak, and we used a web-based survey method to avoid possible infections, causing the sampling of our study was voluntary and conducted by online system. Therefore, the possibility of selection bias should be considered. Third, due to the sudden occurrence of the disaster, we were unable to assess an individual's psychological conditions before the outbreak.

5. Conclusion

In conclusion, we identified a major mental health burden of the Chinese public during the COVID-19 outbreak, and younger people, people spending too much time thinking about the outbreak, and healthcare workers were at a high risk of displaying psychological issues. Previously, when SARS occurred in China, the awareness regarding public's mental health related to the epidemic was low, and no targeted psychological guidelines was available to the public, which was in need during the pandemic period. Therefore, ongoing surveillance and monitoring of the psychological consequences for outbreaks of epidemic-potential, life-threatening diseases, establishing early targeted mental health interventions, should become routine as part of preparedness efforts worldwide.

Author statement contributors

Ning Zhao conceptualized and designed the study, review and revised the manuscript, and approved the final manuscript as submitted. Yeen Huang designed the data collection instruments, coordinated and supervised data collection, carried out the initial analyses, and interpreted the data, drafted the initial manuscript, and approved the final manuscript as submitted. Ning Zhao and Yeen Huang agree to be accountable for all aspects of the study.

Role of the funding source

This work was supported by the National Natural Science Foundation of China. The funder had no role in the design and conduct of the study; management, collection, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Declaration of competing interest

None.

Acknowledgements

The authors would like to thank all the participants in our study. In addition, we express our heartfelt respect to all healthcare workers who are fighting the COVID-19 pandemic on the front line. Finally, we thank Ms. Qiaohong Chen for providing professional language help.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at [doi:10.1016/j.psychres.2020.112954](https://doi.org/10.1016/j.psychres.2020.112954).

References

- Buyse, D.J., Reynolds, C.R., Monk, T.H., Berman, S.R., Kupfer, D.J., 1989. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res* 28, 193–213. [https://doi.org/10.1016/0165-1781\(89\)90047-4](https://doi.org/10.1016/0165-1781(89)90047-4).
- Furer, P., Walker, J.R., Chartier, M.J., Stein, M.B., 1997. Hypochondriacal concerns and somatization in panic disorder. *Depress Anxiety* 6, 78–85 doi: 10.1002/(sici)1520-6394(1997)6:2<78::aid-da4>3.0.co;2-1.
- Gao, W., Ping, S., Liu, X., 2020. Gender differences in depression, anxiety, and stress among college students: a longitudinal study from China. *J Affect Disord* 263, 292–300. <https://doi.org/10.1016/j.jad.2019.11.121>.
- Guo, X., Meng, Z., Huang, G., Fan, J., Zhou, W., Ling, W., Jiang, J., Long, J., Su, L., 2016. Meta-analysis of the prevalence of anxiety disorders in mainland China from 2000 to 2015. *Sci Rep* 6, 28033. <https://doi.org/10.1038/srep28033>.
- Grein, T.W., Kamara, K.B.O., Rodier, G., Plant, A.J., Bovier, P., Ryan, M.J., Ohyama, T., Heymann, D.L., 2000. Rumors of disease in the global village: Outbreak verification. *Emerg Infect Dis* 6, 97–102. <https://doi.org/10.3201/eid0602.000201>.
- Kobayashi, I., Boats, J.M., Delahanty, D.L., 2007. Polysomnographically measured sleep abnormalities in PTSD: a meta-analytic review. *Psychophysiology* 44, 660–669. <https://doi.org/10.1111/j.1469-8986.2007.537.x>.
- Lee, A.M., Wong, J.G., McAlonan, G.M., Cheung, V., Cheung, C., Sham, P.C., Chu, C.M.,

- Wong, P.C., Tsang, K.W., Chua, S.E., 2007. Stress and psychological distress among SARS survivors 1 year after the outbreak. *Can J Psychiatry* 52, 233–240. <https://doi.org/10.1177/070674370705200405>.
- Lee, S.M., Kang, W.S., Cho, A.R., Kim, T., Park, J.K., 2018. Psychological impact of the 2015 MERS outbreak on hospital workers and quarantined hemodialysis patients. *Compr Psychiatry* 87, 123–127. <https://doi.org/10.1016/j.comppsy.2018.10.003>.
- Liu, X.C., Tang, M.Q., Hu, L., Wang, A.Y., Wu, H.X., Zhao, G.F., Gao, C.N., Li, W.S., 1996. Reliability and Validity of Pittsburgh Sleep Quality Index. *J Chin Psychiatry* 29, 103–107. <https://doi.org/10.1007/BF02951625>.
- Lu, Y.C., Shu, B.C., Chang, Y.Y., Lung, F.W., 2006. The mental health of hospital workers dealing with severe acute respiratory syndrome. *Psychother Psychosom* 75, 370–375. <https://doi.org/10.1159/000095443>.
- Mak, I.W., Chu, M.C., Pan, P.C., Yiu, M.G., Chan, V.L., 2009. Long-term psychiatric morbidities among SARS survivors. *Gen Hosp Psychiatry* 31, 318–326. <https://doi.org/10.1016/j.genhosppsych.2009.03.001>.
- Maunder, R., Hunter, J., Vincent, L., Bennett, J., Peladeau, N., Leszcz, M., Sadavoy, J., Verhaeghe, L.M., Steinberg, R., Mazzulli, T., 2003. The immediate psychological and occupational impact of the 2003 SARS outbreak in a teaching hospital. *CMAJ* 168, 1245–1251. <https://doi.org/10.1001/jama.289.18.2432>.
- McAlonan, G.M., Lee, A.M., Cheung, V., Cheung, C., Tsang, K.W., Sham, P.C., Chua, S.E., Wong, J.G., 2007. Immediate and sustained psychological impact of an emerging infectious disease outbreak on health care workers. *Can J Psychiatry* 52, 241–247. <https://doi.org/10.1177/070674370705200406>.
- National Health Commission of the People's Republic of China, 2020. News: 21,569 medical staff have been sent to support Hubei. Accessed February 17, 2020. <http://www.nhc.gov.cn/wjw/xinw/xwzx.shtml>.
- Paules, C.I., Marston, H.D., Fauci, A.S., 2020. Coronavirus infections—more than just the common cold. *JAMA*. Published online January 23, 2020. doi:10.1001/jama.2020.0757.
- People's daily of China, 2020. The novel coronavirus can be inhibited by Chinese patent medicine Shuanghuanglian oral liquid. Accessed January 31, 2020. <http://search.people.com.cn/cnpeople/news/getNewsResult.jsp>.
- Radloff, L.S., 1977. The CES-D scale: A self-report depression scale for research in the general population. *Appl Psychol Meas* 1, 385–401. <https://doi.org/10.1177/014662167700100306>.
- Spitzer, R.L., Kroenke, K., Williams, J.B., Lowe, B., 2006. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med* 166, 1092–1097. <https://doi.org/10.1001/archinte.166.10.1092>.
- Su, T.P., Lien, T.C., Yang, C.Y., Su, Y.L., Wang, J.H., Tsai, S.L., Yin, J.C., 2007. Prevalence of psychiatric morbidity and psychological adaptation of the nurses in a structured SARS caring unit during outbreak: a prospective and periodic assessment study in Taiwan. *J Psychiatr Res* 41, 119–130. <https://doi.org/10.1016/j.jpsychires.2005.12.006>.
- The Japan Times, 2020. Japanese official looking after Wuhan returnees found dead. Accessed February 3, 2020. <https://www.japantimes.co.jp/news/2020/02/02/national/crime-legal/japanese-official-looking-wuhan-returnees-found-dead/>.
- Tong, X., An, D., McGonigal, A., Park, S.P., Zhou, D., 2016. Validation of the Generalized Anxiety Disorder-7 (GAD-7) among Chinese people with epilepsy. *Epilepsy Res* 120, 31–36. <https://doi.org/10.1016/j.eplepsyres.2015.11.019>.
- Wang, D., Hu, B., Hu, C., Zhu, F., Liu, X., Zhang, J., Wang, B., Xiang, H., Cheng, Z., Xiong, Y., Zhao, Y., Li, Y., Wang, X., Peng, Z., 2020. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA*. <https://doi.org/10.1001/jama.2020.1585>. Published online February 7, 2020.
- World Health Organization, 2020a. Novel coronavirus (2019-nCoV): situation report. Accessed February 15, 2020. <https://www.who.int/docs/default-source/coronaviruse/situationreports>.
- World Health Organization, 2020b. Current novel coronavirus (2019-nCoV) outbreak. Accessed January 30, 2020. <https://www.who.int/health-topics/coronavirus>.
- Wang, Y., Chen, R., Zhang, L., 2018. Reliability and validity of generalized anxiety scale-7 in inpatients in Chinese general hospital. *J Clin Psychiatr* 28, 168–171. <https://doi.org/10.3969/j.issn.1005-3220.2018.03.007>.
- Wuhan Municipal Health Commission, 2020. Report of novel coronavirus-infected pneumonia in China. Published January 20, 2020. Accessed January 31, 2020. <http://wjw.wuhan.gov.cn/front/web/showDetail/2020012009077>.
- XINHUANET, 2020. The Latest: China confirms human-to-human transmission of 2019-nCoV, infections among medical staff. Accessed January 21, 2020. http://www.xinhuanet.com/english/2020-01/21/c_138722888.htm.
- Zhang, B.S., Li, J., 2011. Reliability and validity of the simplified version of the Center for Epidemiology Scale for Depression in the national adult population. *Chin Ment Health J* 25, 506–511. <https://doi.org/10.3969/j.issn.1000-6729.2011.07.007>.
- Zhang, J., Wu, Z.Y., Fang, G., Li, J., Han, B.X., Chen, Z.Y., 2010. Establishing a national urban norm for the Center for Epidemiology Scale for Depression. *Chin Ment Health J* 24, 139–143. <https://doi.org/10.3969/j.issn.1000-6729.2010.02.015>.
- Zhu, N., Zhang, D., Wang, W., Li, X., Yang, B., Song, J., Zhao, X., Huang, B., Shi, W., Lu, R., Niu, P., Zhan, F., Ma, X., Wang, D., Xu, W., Wu, G., Gao, G.F., Tan, W., 2020. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med*. <https://doi.org/10.1056/NEJMoa2001017>. Published online January 24, 2020.