

**ERRATUM**

The editorial committee feels sorry for some mistakes in the article “Provision of nutrients enhances Cr<sup>+6</sup> reduction potential of microbial community indigenous to tanneries effluents” on pages 118, 119, 121 and 122 in the previous issue [*Punjab. Univ. J. Zool.*, 20 (1): 113-128]. These pages are being reproduced with corrections in the current issue [Volume 20 (2)] of the journal.

**Table 3: Effect of Cr<sup>6+</sup>, molasses and peptone on number of flagellate inhabitants of original tanneries effluents.**

Experimental Group	Number of flagellates at different days (x10 <sup>9</sup> /ml)								
	1	2	3	4	5	6	7	8	
Control	A <sup>a</sup>	7 <sup>d</sup>	5±1.01	5.67±0.88	2.84±0.60	10.33±0.87	37.83±1.26	42.67±0.76	2.33±0.76
	B <sup>b</sup>	4±0.29 <sup>c</sup>	8.67±0.73	11.34±1.04**	20.5±1.27***	26.5±1.8**	74.5±1.14***	78.33±2.02***	42.84±1.36***
Cr+6 100 µg/ml	A	?	4.83±0.6	5.3±0.33	4.5±0.58	24.67±3.09	15.67±2.83	3.10±3.01	34.65±2.08
	B	3.66±0.6	6±0.77	5.6±0.58	8.84±0.44***	27.17±3.84*	50.17±2.71***	46.67±4.27	45.33±1.91*
Cr+6 200 µg/ml	A	?	2.17±0.33	1.01±0.29	0.5±0.29	1.0±0.29	NVO	NVO	NVO
	B	2±0.29	2.67±0.33	1.67±0.73	2±0.29*	NVO	NVO	NVO	NVO

Each value in table is given as Mean ± SEM of 3 replicates; a: original sample; b: original sample + molasses & peptone; NVO: viable organisms were not detected; \* = P<0.05; \*\* = P<0.01; \*\*\* = P<0.001 (Student's t-test).

**Table 4: Effect of Cr<sup>+6</sup> molasses and peptone on cell volume of flagellate inhabitants of original tanneries effluents.**

Experimental Group	Number of flagellates at different days (x10 <sup>3</sup> /ml)								
	1	2	3	4	5	6	7	8	
Control	A <sup>a</sup>	13.35±0.06	12.6±0.03	12.1±0.06	16.6±0.03	14.99±0.03	9.3±0.006	10.7±0.04	
	B <sup>b</sup>	6.5±0	14.2±0.23*	13.8±0.03***	11.45±0.03***	18.96±0.05***	16±0***	18.6±0.07***	14.44±0***
Cr+6 100 µg/ml	A	?	15.2±0.12	8.6±0.21	9.9±0.16	12.6±0.14	10.33±0.07	8.5±0.05	6.93±0.03
	B	14.4±0.03	10.08±0***	10.3±0***	11.2±0.04***	12.3±0.03*	7.6±0.03***	4.26±0.01	4.3±0.02***
Cr+6 200 µg/ml	A	NVO	8.8±0	8.04±0	6.5±0	8.4±0.1	NVO	NVO	NVO
	B	7.98±0	12.5±0.03***	12.8±0.04***	9.3±0.07***	NVO	NVO	NVO	NVO

Each value in table is given as Mean ± SEM of 3 replicates; a: original sample; b: original sample + molasses & peptone; NVO: viable organisms were not detected, \* = P<0.05; \*\* = P<0.01; \*\*\* = P<0.001 (Student's t-test).

**Table 3: Effect of Cr<sup>6+</sup>, molasses and peptone on number of flagellate inhabitants of original tanneries effluents.**

Experimental Group	Number of flagellates at different days (x10 <sup>3</sup> /ml)								
	1	2	3	4	5	6	7	8	
Control	A <sup>a</sup>	9 <sup>a</sup>	5.67±0.88	2.84±0.60	10.33±0.87	37.83±1.26	42.67±0.76	2.33±0.76	
	B <sup>b</sup>	4±0.29 <sup>c</sup>	8.67±0.73	11.34±1.04*	20.5±1.27***	26.5±1.8**	74.5±1.14**	78.33±2.02***	42.84±1.36***
Cr+6 100 µg/ml	A	?	4.83±0.6	5.3±0.33	4.5±0.58	24.67±3.09	15.67±2.83	3.10±3.01	34.65±2.08
	B	3.66±0.6	6±0.77	5.6±0.58	8.84±0.44***	27.17±3.84*	50.17±2.71***	46.67±4.27	45.33±1.91*
Cr+6 200 µg/ml	A	?	2.17±0.33	1.01±0.29	0.5±0.29	1.0±0.29	NVO	NVO	NVO
	B	2±0.29	2.67±0.33	1.67±0.73	2±0.29*	NVO	NVO	NVO	NVO

Each value in table is given as Mean ± SEM of 3 replicates; a: original sample; b: original sample + molasses & peptone; NVO: viable organisms were not detected. \* = P<0.05; \*\* = P<0.01; \*\*\* = P<0.001 (Student's t-test).

**Table 4: Effect of Cr<sup>+6</sup> molasses and peptone on cell volume of flagellate inhabitants of original tanneries effluents.**

Experimental Group	Number of flagellates at different days (x10 <sup>3</sup> /ml)								
	1	2	3	4	5	6	7	8	
Control	A <sup>a</sup>	13.35±0.06	12.6±0.03	12.1±0.06	16.6±0.03	14.99±0.03	9.3±0.006	10.7±0.04	
	B <sup>b</sup>	6.5±0	14.2±0.23*	13.8±0.03***	11.45±0.03***	18.96±0.05***	16±0***	18.6±0.07***	14.44±0***
Cr+6 100 µg/ml	A	?	15.2±0.12	8.6±0.21	9.9±0.16	12.6±0.14	10.33±0.07	6.93±0.03	
	B	14.4±0.03	10.08±0***	10.3±0***	11.2±0.04***	12.3±0.03*	7.6±0.03***	4.26±0.01	4.3±0.02***
Cr+6 200 µg/ml	A	NVO	8.8±0	8.04±0	6.5±0	8.4±0.1	NVO	NVO	NVO
	B	7.98±0	12.5±0.03***	12.8±0.04***	9.3±0.07***	NVO	NVO	NVO	NVO

Each value in table is given as Mean ± SEM of 3 replicates; a: original sample; b: original sample + molasses & peptone; NVO: viable organisms were not detected, \* = P<0.05; \*\* = P<0.01; \*\*\* = P<0.001 (Student's t-test).

